

Rectifications of Convex Polyhedra

Samuel Reid^{*,†}

April 5, 2016

Abstract

A convex polyhedron, that is, a compact subset of $\mathbb{R}^3$ which is the intersection of finitely many closed half-spaces, can be rectified by taking the convex hull of the mid-points of the edges of the polyhedron. We derive expressions for the side lengths and areas of rectifications of regular polygons in plane, and use these results to compute surface areas and volumes of various convex polyhedra. We introduce rectification sequences, show that there are exactly two disjoint pure rectification sequences generated by the platonic solids, and formulate new results related to the Mahler conjecture.

1 Rectifications of Convex Polygons

We iterate the geometric process of rectification, or maximal truncation, on a seed polyhedron to obtain metric information regarding operations of polyhedra. We introduce the notion of rectification for polygons and polyhedra; rectifications of convex d -polytopes for dimensions $d \geq 4$ will be presented in future work. This topic leads off of the classical polyhedral geometry seen in the works of H.S.M. Coxeter [1],[2], Norman Johnson [3],[4], and others such as Cromwell [5], Conway [6], and Tóth [7]. Intuitively known since antiquity, define:

Definition 1. Let $P_n = \text{conv}\{x_1, \dots, x_n\} \subset \mathbb{R}^2$ be a convex polygon with n vertices and n edges. Then,

$$R_1[P_n] = \text{conv} \left\{ \frac{x_i + x_{i+1}}{2} \mid (x_i, x_{i+1}) \in E(G_{P_n}), 1 \leq i \leq n, x_{n+1} = x_1 \right\},$$

where $E(G_{P_n})$ is the edge set of the graph G_{P_n} ; the combinatorial structure of P_n .

Theorem 1. Let P_n denote a regular n -gon with $\text{area}(P_n) = 1$. Then,

$$\text{area}(R_k(P_n)) = \left(\frac{1 - \cos(\theta_n)}{2} \right)^k$$

^{*}Geometric Energy Corporation. sam@geometricenergy.ca

[†]University of Calgary, Department of Mathematics & Statistics.




Figure 1: The infinite family of polygon rectifications $R_\infty(P_3)$, $R_\infty(P_4)$, and $R_\infty(P_5)$.

and

$$\text{area}(R_\infty(P_n)) = \frac{2}{1 + \cos(\theta_n)}$$

where $\theta_n = \frac{\pi(n-2)}{n}$ is the interior angle of $P_n = R_0(P_n)$, $R_k(P_n)$ is the k^{th} rectification of P_n and

$$R_\infty(P_n) = \bigcup_{k=0}^{\infty} R_k(P_n)$$

Proof. Let P_n be a regular n -gon with $\text{area}(P_n) = 1$. Then by $\text{area}(P_n) = \frac{1}{4}ns^2 \cot\left(\frac{\pi}{n}\right)$, where s is the side length of P_n ,

$$s^2 = \frac{4 \tan\left(\frac{\pi}{n}\right)}{n}$$

By the law of cosines,

$$s_1^2 = \left(\frac{s}{2}\right)^2 + \left(\frac{s}{2}\right)^2 - 2\left(\frac{s}{2}\right)^2 \cos(\theta_n) = \frac{s^2}{2}(1 - \cos(\theta_n)) = \frac{2 \tan\left(\frac{\pi}{n}\right)}{n}(1 - \cos(\theta_n))$$

where $\theta_n = \frac{\pi(n-2)}{n}$ is the interior angle of P_n and s_1 is the side length of $R_1(P_n)$. We can then compute the area of the first rectification of P_n as follows,

$$\text{area}(R_1(P_n)) = \frac{1}{4}ns_1^2 \cot\left(\frac{\pi}{n}\right) = \frac{1 - \cos(\theta_n)}{2}$$

Now, let $k \geq 2$ be arbitrary and let s_k be the side length of $R_k(P_n)$. For induction, assume that

$$s_k^2 = \frac{\tan\left(\frac{\pi}{n}\right)}{2^{k-2}n}(1 - \cos(\theta_n))^k$$

By the law of cosines,

$$s_{k+1}^2 = \frac{s_k^2}{2}(1 - \cos(\theta_n)) = \frac{\tan\left(\frac{\pi}{n}\right)}{2^{k-1}n}(1 - \cos(\theta_n))^{k+1}$$

Then,

$$\text{area}(R_k(P_n)) = \frac{1}{4}ns_k^2 \cot\left(\frac{\pi}{n}\right) = \left(\frac{1 - \cos(\theta_n)}{2}\right)^k$$

implies that

$$\text{area}(R_{k+1}(P_n)) = \frac{1}{4}ns_{k+1}^2 \cot\left(\frac{\pi}{n}\right) = \left(\frac{1 - \cos(\theta_n)}{2}\right)^{k+1}$$

Therefore the formula for $\text{area}(R_k(P_n))$ holds for all $k \in \mathbb{N}$ by induction. We now compute the area of the infinite family of polygon rectifications $R_\infty(P_n)$.

$$\text{area}(R_\infty(P_n)) = \text{area}\left(\bigcup_{k=0}^{\infty} R_k(P_n)\right) = \sum_{k=0}^{\infty} \text{area}(R_k(P_n)) = \sum_{k=0}^{\infty} \left(\frac{1 - \cos(\theta_n)}{2}\right)^k = \frac{2}{1 + \cos(\theta_n)}$$

by the convergence of the geometric series since $\left|\frac{1 - \cos(\theta_n)}{2}\right| < 1$ for all P_n with $n \geq 3$. $\square$

While the perimeter of the infinite rectification $R_\infty(P_n)$ follows trivially from the proof of Theorem 1, it is in general a difficult problem to determine the surface area of $R_\infty(\text{conv}\{P\})$ when $P \subset \mathbb{R}^d$ with $d \geq 3$.

Corollary 1. *Let P_n be a regular n -gon with $\text{area}(P_n) = 1$. Then,*

$$\text{Perimeter}(R_\infty(P_n)) = \frac{2\sqrt{n \tan(\pi/n)}}{1 - |\cos(\pi/n)|}$$

Proof. We have by the proof of Theorem 1 that

$$s_k = \sqrt{\frac{\tan\left(\frac{\pi}{n}\right)}{2^{k-2}n}}(1 - \cos(\theta_n))^k$$

thus,

$$\text{Perimeter}(R_\infty(P_n)) = \sum_{k=0}^{\infty} ns_k = 2\sqrt{n \tan\left(\frac{\pi}{n}\right)} \sum_{k=0}^{\infty} \left(\sqrt{\frac{1 - \cos(\theta_n)}{2}}\right)^k = \frac{\sqrt{8n \tan\left(\frac{\pi}{n}\right)}}{\sqrt{2} - \sqrt{1 - \cos(\theta_n)}}$$

by the convergence of the geometric series since $\left|\sqrt{\frac{1 - \cos(\theta_n)}{2}}\right| < 1$ for all P_n with $n \geq 3$. Since $n > 0$ we have by the definition of θ_n that

$$\text{Perimeter}(R_\infty(P_n)) = \frac{2\sqrt{n \tan(\pi/n)}}{1 - |\cos(\pi/n)|}$$

$\square$

We thus have a complete classification of regular polygons rectification sequences and their area and perimeter. We now want to generalize this problem to classifying rectification sequences of polyhedra in $\mathbb{R}^3$ and computing their geometric measures.

2 Rectifications of Convex Polyhedra

Definition 2. Let $P = \text{conv}\{x_1, \dots, x_v\} \subset \mathbb{R}^3$ be a convex polyhedron with e edges and f faces. Then,

$$R_1[P] = \text{conv}\left\{\frac{x_i + x_j}{2} \mid (x_i, x_j) \in E(G_P)\right\},$$

where $E(G_P)$ is the edge set of the graph G_P ; the combinatorial structure of P .

Rectifications can be thought of as the maximal truncation of a polyhedron which sends a vertex to a face. If we consider the truncation of each vertex of a polyhedron by a plane with a normal vector coinciding with the centroid ray of the polyhedron then we obtain the rectification of the polyhedron when each new edge created by the truncation touches another newly created edge.




Figure 2: The first two rectifications of a tetrahedron, namely an octahedron and a cuboctahedron inscribed inside each other.

Rectifications of convex polyhedra can be easily understood combinatorially due to $V - E + F = \chi = 2$, as the rectification of a polyhedron can be thought of as a map between f -vectors

$$(v, e, f) \mapsto (e, 2e, 2 + e)$$

There are two distinct infinite sequences of polytope rectifications which include the platonic solids; the first, denoted by η_s includes the tetrahedron, octahedron, and cube and has an f -vector sequence given by

$$(4, 6, 4) \xrightarrow{\text{Tetrahedron}} (6, 12, 8) \xrightarrow{\text{Octahedron}} (12, 24, 14) \xrightarrow{\text{Cuboctahedron}} \dots$$

and the second, denoted by ξ_s includes the icosahedron and dodecahedron, which have the property that they have equal rectifications, and has an f -vector sequence given by

$$(12, 30, 20) \xrightarrow{\text{Icosahedron}} (30, 60, 32) \xrightarrow{\text{Icosidodahedron}} (60, 120, 62) \xrightarrow{\text{Icosidodecahedron}} \dots$$

Theorem 2. *The polytope rectification sequences η_s and ξ_s are disjoint.*

Proof. By the combinatorial characterization of rectification as an f -vector map $(v, e, f) \mapsto (e, 2e, 2 + e)$, an equivalent question is whether there exists a polytope in the sequence η_s which also exists in ξ_s , which implies that the existence of a pair (k, x) which satisfies

$$2^k 12 = 2^x 30$$

but this is never satisfied for $k, x \in \mathbb{Z}$ as $k - x = \frac{\log(5)}{\log(2)} - 1 \notin \mathbb{Q}$. □




Figure 3: The first rectification of the pentaprism.

We now calculate the volume of the first rectification of a prism generated from a regular polygon in the plane.

Theorem 3. *Let P_n be a regular polygon of area 1 in the plane. Then,*

$$\text{vol}(R_1(P_n \times I)) = 1 - \frac{s^2}{48} \sin(\theta_n)$$

where s is the side length of P_n and $\theta_n = \frac{\pi(n-2)}{n}$ is the internal angle of a regular n -gon.

3 Pure Rectification Sequences

Definition 3. A pure rectification sequence of length k is a sequence

$$P \rightarrow R_1[P] \rightarrow \cdots \rightarrow R_{k-1}[P]$$

so that every element of the sequence is semiregular.

Theorem 4. Let

$$P \rightarrow R_1[P] \rightarrow \cdots \rightarrow R_{k-1}[P]$$

be a semiregular rectification sequence. Then the f -vector and volume of $R_k[P]$ are

$$(f_0(R_k[P]), f_1(R_k[P]), f_2(R_k[P])) = (2^{k-1}f_1(P), 2^k f_1(P), 2 + 2^{k-1}f_1(P))$$

and

$$\text{vol}(R_k[P]) = \text{vol}(P) - \sum_{i=1}^{f_0(P)} \text{vol}(\text{conv}\{P/v_i^0 \cup \{v_i^0\}\}) - \sum_{N=1}^k \sum_{j=1}^{2^{N-1}f_1(P)} \text{vol}(\text{conv}\{R_N[P]/v_j^N \cup \{v_j^N\}\})$$

where P/v_i^0 denotes the vertex figure of P at $v_i^0 \in V(P)$ truncated to half the edge length of P and $R_N[P]/v_j^N$ denotes the vertex figure of $R_N[P]$ at the vertex $v_j^N \in V(R_N[P])$ truncated to half the edge length of $R_N[P]$.

Conjecture 1. Let P be a convex polyhedron. If $R_1[P]$ is contained in a pure rectification sequence then

$$R_1[P] = \lambda R_1[P^\circ], \exists \lambda > 0.$$

Lemma 1. Let P be a convex polyhedron so that $R_1[P] = \lambda R_1[P^\circ], \exists \lambda > 0$. If

$$\text{vol}(R_1[P]) \geq \sqrt{\frac{32\lambda}{3}}$$

then the Mahler conjecture holds for $R_1[P]$ and $R_1[P] \times R_1[P]^\circ$.

Conjecture 2 (Mahler Conjecture for Pure Rectifications and a Class of Hanner 6-Polytopes). If $R_1[P]$ is a pure rectification of a convex polyhedron P then the Mahler conjecture is true for $R_1[P]$ and $R_1[P] \times R_1[P]^\circ$.

Proof. Apply Conjecture 1 and Lemma 1. □

Conjecture 3. Let $P \subset \mathbb{R}^3$ be a convex polyhedron. Then,

$$\exists \lambda \in \mathbb{R}^+ : \lambda P^\circ \cap P = R_1[P].$$

References

- [1] H.S.M. Coxeter, M.S. Longuet-Higgins, J.C.P. Miller. Uniform Polyhedra. 1954, Philosophical Transactions of the Royal Society of London 246 A (1954), pp. 401450.
- [2] H.S.M. Coxeter. Regular Polytopes. 1973, Dover.
- [3] N. Johnson. Convex polyhedra with regular faces. 1966, Canadian Journal of Mathematics 18: 169200.
- [4] N. Johnson. The Theory of Uniform Polytopes and Honeycombs. 1966, Ph.D. Dissertation, University of Toronto.
- [5] P. Cromwell. Polyhedra. 1977, Cambridge University Press.
- [6] John H. Conway, Heidi Burgiel, Chaim Goodman-Strass. The Symmetries of Things. 2008, CRC Press.
- [7] László Fejes Tóth. Regular Figures. 1964, Oxford University Press.